

LEAFLET: INFORMATION FOR THE USER

Freshalgin 500th mg tablets

Metamizole sodium (*M etamizol sodium*)

Read this leaflet carefully before you start taking this medicine.

This medicine is available without prescription. However, you have to follow instructions to get the best results from treatment. Keep this leaflet. You may need to read it again.

If you have any further questions, ask your doctor or pharmacist.

If any of the side effects gets serious, or you notice other effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Freshalgin and what it is used
2. Before taking Freshalgin
3. How to take Freshalgin
4. Possible side effects
5. How to store Freshalgin
6. Additional Information

1. WHAT AND FRESHALGIN WHAT IT IS USED

Freshalgin non-opioid medicine is a strong analgesic effect and lower fever and moderate inflammatory ystvie December.

Freshalgin shown:

- for the short chno treatment of pain of various origins and intensity when other agents are ineffective or contraindicated and our headache, toothache, joint and muscle pain and inflammation, arthritis, neuralgia, post-traumatic and postoperative pain in, pain and cramps etc and renal s roads and bile.
- To relieve pain in cancer patients, such as "penultimate stage" prior to narcotic analgesics.
- Short lowering high body temperature which is not affected by other medicines or if they are contraindicated.

2. BEFORE YOU TAKE FRESHALGIN

Do not take this medicine if:

- If you are vrahchuvstviteln and the M e tamizol active substance and / or any of the excipients;
- Hypersensitivity to pirazonovi derivatives or other NSAIDs;
- Genetic deficiency of glucose-6-phosphate dehydrogenase;
- Bronchial asthma;
- Hematologic Diseases - leukopenia (reduction in the total number of leukocytes), agranulocytosis, hemolytic anemia, hepatic porphyria, aplastic anemia (anemia due udrezhdane haematopoietic bone marrow function);
- Severe liver or kidney disease;
- Pregnancy and lactation
- Children under 7 years

Take special care with Freshalgin

- In patients with renal function in damages, gastric and duodenal yazv and only be used under medical supervision.
- Diseases occurring by reducing the number of leukocytes requires tyahnot o tracking to reduce the risk of leucopenia.
- When driven allergic asthma.
- With use urine may turn red.

Taking other medicines

Please yte inform your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Concomitant use of Freshalgin other medicines can strengthen or weaken their effects:

- M e tamizol can reduce the activity of coumarin anticoagulants, resulting in accelerating their degradation (metabolism);
- Coadministration of M e tamizol drug, accelerating its degradation called "enzyme inducers" of liver metabolism (Barbiturates, glutethimide, phenylbutazone) can reduce its effects;
- Tricyclic antidepressants, oral contraceptives and allopurinol, enhance the effects of M e tamizol due to slow its metabolism;
- M e tamizol lowered plasma levels of Cyclosporine with concomitant use.
- The simultaneous administration of Metamizol with chlorpromazine and other phenol derivatives iazinovi an increased risk of lower body temperatura;
- Co-administration of M e tamizol other antipyretic, analgesic and anti-inflammatory drugs increase the risk of allergic reactions;
- There is a risk of suppression of bone marrow function In association with chloramphenicol and other myelotoxic products.

Taking Freshalgin with food and drink

A alcohol seen to strengthen the analgesic effect of Metamizol.

Pregnancy

Please consult your doctor or advice before taking any medicine.

Freshalgin not recommended p o pregnancy.

Breastfeeding

The product passes into breast milk and is not recommended during lactation.

Driving and operating machinery

Product not affect your ability to drive or operate machinery.

Important information nyak OI component of Freshalgin

This product contain as an ingredient excipient wheat starch, which makes it unsuitable for patients intolerant to gluten.

3. HOW TO TAKE FRESHALGIN

Adults and children 15 and above years, without prescription:

Therapeutic dose - 1 tablet 3 to 4 times a day but.

Maximum single dose of 2 tablets (1,0 g)

Maximum daily dose 6 tablets (3, 0 g)

Children and adolescents under 15 years of age (body weight <53 kg) - on prescription!

Therapeutic doses:

Children from 7-9 years-½ tablet 2 times daily, maximum daily dose of 2 tablets.

Children 10-14 years - ½ 1 tablet to 3 times daily, maximum daily dose of 3 tablets.

Elderly

In the elderly should be given to reducing the dose to the possibility of decreased elimination of the active substance or its metabolites.

In patients with impaired renal or hepatic function, the dose of metamizole sodium should be reduced.

It is recommended Freshalgin be taken after meals, if necessary, before meals, with plenty of fluids.

The duration of the treatment course without consultation with a doctor should not exceed 5 days in adults and in children 3 days.

If symptoms persist, consult your doctor.

If you take a large dose Freshalgin than prescribed

When taking a higher dose than prescribed, you should consult a doctor for help!

In case of overdose can occur following symptoms: nausea, vomiting, dizziness, tinnitus, decrease in body temperature, reduced urine output, bluish skin, shortness of breath. It has very low blood pressure to shock, acute liver failure, tonic-clonic convulsions, coma.

The first steps in overdose are inducing vomiting, stomach lavage old, administration of activated charcoal, followed by symptomatic treatment. There is no specific antidote.

If you miss me, they adopted Freshalgin

If you miss a dose, take the medicine at the next regular intake without increasing the dose.

4. POSSIBLE SIDE EFFECTS

Like all medicines and Freshalgin can cause side effects. Most commonly seen in hypersensitivity to the drug.

Can be observed

- Immune system - allergies in very rare cases, allergic shock;
- Nervous system - with high doses of epileptiform convulsions;
- Sense organs - dizziness;
- Cardiovascular system - a rare feeling of palpitations, rapid pulse, cyanosis, low blood pressure;
- Blood and lymphatic system - rare low red and white blood cells and platelets, agranulocytosis (Reduction or lack of certain white blood cells), hemolytic anemia, leukopenia (low leukocytes) thrombocytopenia (reduced platelet count);
- Respiratory System - sore throat, shortness of breath frequently;
- Digestive system - Loss of appetite, nausea, vomiting, retention of bile, jaundice.
- Genitourinary system - with high doses of renal impairment;
- Cutaneous allergic reactions, rash, urticaria, edema Kvinkve, itching, rarely break the surface layer of the skin, Stevens - Johnson syndrome or Lyle.

If any of these side effects gets serious, or if you are not mentioned in this leaflet, please tell your doctor or pharmacist.

5. STORAGE FRESHALGIN

Store protected from light below 25 ° C.

Keep your tablets in the original container.

Keep out of reach of children!

Do not use after the expiry date printed on the package. Expiry date refers to the last day of that month.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. ADDITIONAL INFORMATION

What Freshalgin

- The active substance is sodium metamizole. Each tablet contains 500 mg of sodium metamizole (*Metamizole sodium*).

- Excipients wheat starch, gelatin, povidone, microcrystalline cellulose, magnesium stearate and talc.

Freshalgin looks like and contents of pack

The tablets are white or off-white, round, flat, Scored on one side. The tablet can be divided into two equal halves.

10 tablets in blister PVC / A 1.

1 or 2 blisters with one sheet per carton.

Marketing Authorisation Holder and Manufacturer

Adipharm Ltd.

Boulevard Vitosha № 130

1700 Sofia, Bulgaria